

**Boris Baeumer · Mihály Kovács · Mark M.
Meerschaert**

Fractional reproduction-dispersal equations and heavy tail dispersal kernels

May 30, 2007

Abstract Reproduction-Dispersal equations, called reaction - diffusion equations in the physics literature, model the growth and spreading of biological species. Integro-Difference equations were introduced to address the shortcomings of this model, since the dispersal of invasive species is often more widespread than what the classical RD model predicts. In this paper, we extend the RD model, replacing the classical second derivative dispersal term by a fractional derivative of order $1 < \alpha \leq 2$. Fractional derivative models are used in physics to model anomalous super-diffusion, where a cloud of particles spreads faster than the classical diffusion model predicts. This paper also establishes a connection between the new

Partially supported by the Marsden fund administered by the Royal Society of New Zealand, NSF grants DMS-0139927, DMS-0417869 and DMS-0706440, and Postdoctoral grant No. 623-2005-5078 of the Swedish Research Council.

Boris Baeumer
Department of Mathematics and Statistics, University of Otago,
P.O. Box 56, Dunedin 9001, New Zealand
Tel.: +643-479-7763
Fax: +643-479-8427
E-mail: bbaeumer@maths.otago.ac.nz

Mihály Kovács
Department of Mathematics and Statistics, University of Otago,
P.O. Box 56, Dunedin 9001, New Zealand
Tel.: +643-479-7889
Fax: +643-479-8427
E-mail: mkovacs@maths.otago.ac.nz

Mark M. Meerschaert
Department of Statistics and Probability
Michigan State University
A413 Wells Hall
East Lansing, MI 48824-1027 USA
Tel.: +1-517-432-7097
Fax: +1-517-432-1405
E-mail: mcubed@stt.msu.edu

RD model and a corresponding ID equation with a heavy tail dispersal kernel. The general theory developed here accommodates a wide variety of infinitely divisible dispersal kernels that adapt to any scale. Each one corresponds to a generalised RD model with a different dispersal operator. The connection established here between RD and ID equations can also be exploited to generate convergent numerical solutions of RD equations along with explicit error bounds.

Keywords reproduction-dispersal equation · integro-difference equation · fractional derivative · anomalous diffusion · operator splitting

1 Introduction

The classical reproduction-dispersal (RD) equation

$$\frac{\partial u}{\partial t} = f(u) + D \frac{\partial^2 u}{\partial x^2}. \quad (1)$$

for the growth and dispersal of biological species [25, 35] can underestimate the speed of invasion. This has led to the consideration of the integro-difference (ID) equation

$$u(x, t + \tau) = \int_{-\infty}^{\infty} k^\tau(x, y) g_\tau(u(y, t)) dy \quad (2)$$

that replaces the classical second derivative dispersal term by a convolution with a dispersal kernel [36, 52, 53]. In both models, the population density is $u(x, t)$ at location x and time t . In the physics literature, (1) is called the reaction-diffusion equation. The ID model with a Gaussian dispersal kernel is essentially equivalent to the classical RD model, in a sense that will be made precise in this paper. Alternative dispersal kernels with heavier tails model the faster and wider spreading observed in many field studies [15, 18, 19, 32, 34, 55]. This paper proposes a new fractional RD equation

$$\frac{\partial u}{\partial t} = f(u) + D \frac{\partial^\alpha u}{\partial |x|^\alpha} \quad (3)$$

with $1 < \alpha \leq 2$, extending the classical approach. Fractional derivatives are used in physics to model anomalous diffusion, where a cloud of particles spreads farther and faster than the classical diffusion model predicts [8, 12, 13, 33, 43–46, 48, 49, 60, 77]. Fractional derivative models have also been proposed in finance [27, 58, 59, 64–66] to model price volatility, and in hydrology [6, 9–11, 67, 68] to model fast spreading of pollutants. In each case, the fractional derivative term substitutes for the classical second derivative term, resulting in a wider and faster spread. Fractional derivatives are a natural analogue of their integer-order cousins [50, 61]. They were invented by Leibnitz, but their utility in modelling practical situations has only recently been recognised. In this paper, we develop a general theory of RD equations including the classical and fractional versions. We also prove an explicit connection between RD and ID models with infinitely divisible dispersal kernels that adapt to any scale. The dispersal kernels that correspond to the fractional RD model are stable probability densities [62] that occur in the generalised central limit theorem of statistics [24, 26]. The connection between ID and generalised RD models also yields numerical solutions with explicit error bounds, based on operator splitting.

2 The general Reproduction-Dispersal equation

The classical reproduction-dispersal equation (1) and its fractional analogue (3) are both special cases of a general form

$$\frac{\partial}{\partial t} u(x, t) = Au(x, t) + f(u(x, t)), \quad u(x, 0) = u_0(x) \quad (4)$$

where A is a pseudo-differential operator [31] that appears as the generator of some continuous convolution semi-group [4, 56]. Our goal in this section is to explore the connection between this continuous time evolution equation and its discrete time analogue, the integro-difference equation

$$u_{n+1}(x) = \int_{-\infty}^{\infty} k^\tau(x, y) g_\tau(u_n(y)) dy \quad (5)$$

where $u_n(x) = u(x, n\tau)$ with some $\tau > 0$ fixed. Our general approach is based on operator theory for abstract differential equations [4, 23, 56] and infinitely divisible probability distributions [24, 47].

Let X be a Banach space of functions $h : \mathbb{R} \rightarrow \mathbb{R}$ with associated norm $\|h\|$, and rewrite the RD equation (4) in operator theory notation as

$$\dot{u}(t) = Au(t) + f(u(t)), \quad t > 0, \quad u(0) = u_0 \quad (6)$$

where $u : [0, \infty) \rightarrow X$ and A is the generator of a strongly continuous semi-group $T(t)$ on X . If $f : X \rightarrow X$ is Lipschitz continuous, then (6) has a unique global mild solution $u(t) := W(t)u_0$ for all $u_0 \in X$ [56, Section 6.1], i.e., u is continuous and satisfies the corresponding integral equation

$$u(t) = T(t)u_0 + \int_0^t T(t-s)f(u(s)) ds. \quad (7)$$

The growth or reproduction equation $\dot{u} = f(u)$ is a special case of (6) with $A = 0$. Denote its unique mild solution by $u(t) := S(t)u_0$. Then u is also a strong solution (see, e.g., [30, page 67]) and the nonlinear operators $S(t)$ form a semi-group called the flow of the abstract differential equation $\dot{u} = f(u)$. The solution operator $W(t)u_0$ to the abstract RD equation (6) can then be computed using the Trotter product formula [14, 20, 51]

$$W(t)u_0 = \lim_{n \rightarrow \infty} [T(\frac{t}{n})S(\frac{t}{n})]^n u_0 = \lim_{n \rightarrow \infty} [S(\frac{t}{n})T(\frac{t}{n})]^n u_0, \quad u_0 \in X. \quad (8)$$

This operator splitting expresses solutions in terms of the two sub-problems, since $T(t)u_0$ solves the dispersion equation $\dot{u} = Au$ and $S(t)u_0$ solves the growth equation $\dot{u} = f(u)$.

Assume that X is an *ordered Banach space*, i.e., a real Banach space endowed with a partial ordering \leq such that

1. $u \leq v$ implies $u + w \leq v + w$ for all $u, v, w \in X$.
2. $u \geq 0$ implies $\lambda u \geq 0$ for all $u \in X$ and $\lambda \geq 0$.
3. $0 \leq u \leq v$ implies $\|u\| \leq \|v\|$ for all $u, v \in X$.
4. The positive cone $X_+ := \{u \in X : u \geq 0\}$ is closed.

A typical example of an ordered Banach space is $C_0(\mathbb{R})$, the space of continuous functions $u : \mathbb{R} \rightarrow \mathbb{R}$ such that $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$, endowed with the supremum norm $\|u\| = \sup\{|u(x)| : x \in \mathbb{R}\}$, and endowed with the partial ordering $u \leq v$ whenever $u(x) \leq v(x)$ for all $x \in \mathbb{R}$. An operator A on an ordered Banach space is positive if $0 \leq u \leq v$ implies $0 \leq Au \leq Av$. We also write $B \leq A$ if $0 \leq Bu \leq Au$ for any $u \geq 0$. The following result will be used to associate each abstract RD equation with an associated ID (integro-difference) equation.

Proposition 1 *Let X be an ordered Banach space, and assume that the strongly continuous semi-group $T(\cdot)$ generated by the linear operator A and the nonlinear semi-group $S(\cdot)$ generated by the Lipschitz continuous function f are positive. If*

$$T(t)S(t)u_0 \leq S(t)T(t)u_0 \quad (9)$$

holds for all $t \in [0, T]$ and $u_0 \geq 0$, then the unique mild solution $W(t)u_0$ of the abstract reaction-diffusion Equation (6) is given by (8) and satisfies

$$\begin{aligned} [T(\frac{t}{n})S(\frac{t}{n})]^n u_0 &\leq [T(\frac{t}{2n})S(\frac{t}{2n})]^{2n} u_0 \leq W(t)u_0 \\ &\leq [S(\frac{t}{2n})T(\frac{t}{2n})]^{2n} u_0 \leq [S(\frac{t}{n})T(\frac{t}{n})]^n u_0 \end{aligned} \quad (10)$$

for all $u_0 \geq 0$, $n \in \mathbb{N}$ and $t \in [0, T]$.

Proof We follow [20, Theorem 15]. It follows from our assumption that $W(t)u_0$ is given by (8). Next we show that

$$\begin{aligned} [T(\frac{t}{n})S(\frac{t}{n})]^n u_0 &\leq [T(\frac{t}{2n})S(\frac{t}{2n})]^{2n} u_0 \\ &\leq [S(\frac{t}{2n})T(\frac{t}{2n})]^{2n} u_0 \leq [S(\frac{t}{n})T(\frac{t}{n})]^n u_0, \end{aligned} \quad (11)$$

$u_0 \geq 0$, $n \in \mathbb{N}$ and $t \in [0, T]$. For $u_0 \geq 0$ using (9) repeatedly we have

$$\begin{aligned} T(\frac{t}{n})S(\frac{t}{n})u_0 &= T(\frac{t}{2n})T(\frac{t}{2n})S(\frac{t}{2n})S(\frac{t}{2n})u_0 \\ &\leq T(\frac{t}{2n})S(\frac{t}{2n})T(\frac{t}{2n})S(\frac{t}{2n})u_0 \leq S(\frac{t}{2n})T(\frac{t}{2n})T(\frac{t}{2n})S(\frac{t}{2n})u_0 \\ &\leq S(\frac{t}{2n})T(\frac{t}{2n})S(\frac{t}{2n})T(\frac{t}{2n})u_0 \leq S(\frac{t}{2n})S(\frac{t}{2n})T(\frac{t}{2n})T(\frac{t}{2n})u_0 \\ &= S(\frac{t}{n})T(\frac{t}{n})u_0. \end{aligned}$$

For any operators $A, B : X \rightarrow X$, it is not hard to check that the inequality $0 \leq B \leq A$ implies $0 \leq B^n \leq A^n$ ($n \in \mathbb{N}$) which finishes the proof of (11). Fix $n \in \mathbb{N}$ and let

$$u_k := \left[T(\frac{t}{n2^k})S(\frac{t}{n2^k}) \right]^{n2^k} u_0.$$

By (11), $u_0 \leq u_1 \leq \dots \leq u_k \leq \dots$ and $u_k \rightarrow W(t)u_0$ as $k \rightarrow \infty$ by (8). Therefore, it follows from the closedness of the positive cone X_+ that $u_k \leq W(t)u_0$ for all $k = 0, 1, \dots$. This shows the second inequality in (10). A similar argument yields the third inequality and the proof is complete.

For a detailed and exhaustive introduction to positive semi-groups we refer to [5]. One useful class of positive strongly continuous semi-groups are the infinitely divisible semi-groups, which are associated with certain families of probability densities. Suppose that Y is a random variable on \mathbb{R} with probability density $k(y)$ and Fourier transform $\hat{k}(\lambda) = \int e^{-i\lambda y} k(y) dy$. Let $k^n = k * \dots * k$ denote the n -fold convolution of k with itself. We say that Y (or k) is *infinitely divisible* if for each $n = 1, 2, 3, \dots$ there exist independent random variables Y_{n1}, \dots, Y_{nm} with the same density k_n such that $Y_{n1} + \dots + Y_{nm}$ is identically distributed with Y . The normal, Cauchy, double-Gamma, Laplace, α -stable, and Student- t densities are all infinitely divisible. The Lévy representation (see, e.g., Theorem 3.1.11 in [47]) states that if k is infinitely divisible then $\hat{k}(\lambda) = e^{\psi(\lambda)}$ where

$$\psi(\lambda) = -i\lambda a - \frac{1}{2}\lambda^2 b^2 + \int_{y \neq 0} \left(e^{-i\lambda y} - 1 + \frac{i\lambda y}{1+y^2} \right) \phi(y) dy, \quad (12)$$

where $a \in \mathbb{R}$, $b \geq 0$, and the *jump intensity* ϕ satisfies

$$\int_{y \neq 0} \min\{1, y^2\} \phi(y) dy < \infty.$$

The unique triple $[a, b, \phi]$ is called the Lévy representation of the infinitely divisible density k . It follows that we can define the convolution power k^t to be the infinitely divisible density with Lévy representation $[ta, tb, t\phi]$, so that k^t has Fourier transform $e^{t\psi(k)}$ for any $t \geq 0$. This simple fact will allow us to define appropriate dispersal kernels at any scale.

Any infinitely divisible density is associated [31, Example 4.1.3] with a strongly continuous semi-group on $C_0(\mathbb{R})$ defined by

$$[T(t)u](x) := \int_{-\infty}^{\infty} k^t(x-y) u(y) dy. \quad (13)$$

Every function $u \in C_0(\mathbb{R})$ with $u', u'' \in C_0(\mathbb{R})$ belongs to the domain of the generator A of the semi-group (13), and for such functions we have [63, Theorem 31.5]

$$\begin{aligned} Au(x) = & -au'(x) + \frac{1}{2}b^2u''(x) \\ & + \int_{y \neq 0} \left(u(x-y) - u(x) + \frac{u'(x)y}{1+y^2} \right) \phi(y) dy. \end{aligned} \quad (14)$$

We show now that in case of Fisher's equation, where the nonlinearity f is not globally Lipschitz, the solution of the discrete ID equation still yields bounds on the solution of the continuous RD equation and the solution of the ID equation converge to the unique solution of the RD equation as the time step tends to zero.

Theorem 1 *Let k be infinitely divisible and let A denote the generator of the associated strongly continuous convolution semi-group $T(t)$ defined by (13) on $X := C_0(\mathbb{R})$. Let $f(u) = ru(1 - u/K)$. Then (6) with initial condition $u_0 \geq 0$ has a unique mild solution $u(t) = W(t)u_0$ for all $u_0 \geq 0$ in X given by the Trotter product formula*

$$W(t)u_0 = \lim_{n \rightarrow \infty} \left[T\left(\frac{t}{n}\right) S\left(\frac{t}{n}\right) \right]^n u_0 = \lim_{n \rightarrow \infty} \left[S\left(\frac{t}{n}\right) T\left(\frac{t}{n}\right) \right]^n u_0. \quad (15)$$

Moreover,

$$\begin{aligned} [T(\frac{t}{n})S(\frac{t}{n})]^n u_0 &\leq [T(\frac{t}{2n})S(\frac{t}{2n})]^{2n} u_0 \leq W(t)u_0 \\ &\leq [S(\frac{t}{2n})T(\frac{t}{2n})]^{2n} u_0 \leq [S(\frac{t}{n})T(\frac{t}{n})]^n u_0 \end{aligned} \quad (16)$$

for all $n \in \mathbb{N}$, where

$$S(t)u_0 = \frac{Ku_0 \exp(rt)}{K + (\exp(rt) - 1)u_0}. \quad (17)$$

Proof As the growth function in Fisher's equation is not Lipschitz, we introduce a cut off function (which is Lipschitz) via the function $\tilde{f}_N : \mathbb{R} \rightarrow \mathbb{R}$ as

$$[f_N(u)](x) = \tilde{f}_N(u(x)) := \begin{cases} 0 & \text{if } u(x) < 0 \\ ru(x) \left(1 - \frac{u(x)}{K}\right) & \text{if } 0 \leq u(x) \leq NK \\ rNK(1 - N) & \text{if } u(x) > NK. \end{cases} \quad (18)$$

Choose $N \geq 2$ such that

$$0 \leq u_0(x) \leq NK \text{ for all } x \in \mathbb{R}. \quad (19)$$

Note that the initial population density $u_0(x)$ may exceed the carrying capacity. We are going to show that the solution to (6) is the same as the solution to the abstract reproduction-dispersal equation

$$\dot{u}(t) = Au(t) + f_N(u(t)), \quad u(0) = u_0 \geq 0. \quad (20)$$

if N is chosen according to (19). As f_N is Lipschitz, the unique mild solution $u_N(t) = W_N(t)u_0$ of (20) is again given by the Trotter product formula

$$\begin{aligned} u_N(t) = W_N(t)u_0 &= \lim_{n \rightarrow \infty} [T(\frac{t}{n})S_N(\frac{t}{n})]^n u_0 \\ &= \lim_{n \rightarrow \infty} [S_N(\frac{t}{n})T(\frac{t}{n})]^n u_0, \end{aligned} \quad (21)$$

where $S_N(\cdot)$ is the nonlinear positive semi-group generated by f_N . The positivity of $S_N(\cdot)$ follows from the uniqueness of solutions of $\dot{u} = f_N(u)$, $u(0) = u_0$, and the fact that $\tilde{f}_N(0) = 0$.

Since k^t is a probability density function, the semi-group $T(\cdot)$ applied to u_0 , with $0 \leq u_0(x) \leq NK$ for all x , satisfies

$$0 \leq [T(t)u_0](x) \leq NK \int_{-\infty}^{\infty} k^t(x-y)dy = NK. \quad (22)$$

Furthermore, [7, Proposition 4.2] shows that

$$0 \leq [S_N(t)u_0](x) = [S(t)u_0](x) \leq NK, \quad x \in \mathbb{R}$$

provided (19) holds. Hence,

$$0 \leq [T(\frac{t}{n})S_N(\frac{t}{n})]^n u_0(x) = [T(\frac{t}{n})S(\frac{t}{n})]^n u_0(x) \leq NK \quad (23)$$

for all $x \in \mathbb{R}$ and

$$0 \leq [S_N(\frac{t}{n})T(\frac{t}{n})]^n u_0(x) = [S(\frac{t}{n})T(\frac{t}{n})]^n u_0(x) \leq NK \quad (24)$$

for all $x \in \mathbb{R}$. This also shows that $0 \leq [u_N(t)](x) \leq NK$ for all $x \in \mathbb{R}$ in view of (21). Since $f_N(u(x)) = f(u(x))$ for $0 \leq u(x) \leq NK$, $u_N(t)$ is a mild solution of (6) as well. Since f is locally Lipschitz, a well known result [56, Chapter 6, Theorem 1.4] implies that (6) has a unique local mild solution and since $u_N(t)$ is defined for all $t > 0$ it follows that $u_N(t)$ is the unique global mild solution of (6) and is given by the Trotter product formula (15).

Finally, an easy computation shows that the function

$$y \mapsto [\tilde{S}(t)](y) := \frac{Ky \exp(rt)}{K + (\exp(rt) - 1)y}$$

is concave down on $y > 0$ for any $t > 0$ and $[S(t)u_0](x) = [\tilde{S}(t)](u_0(x))$. Therefore, by Jensen's inequality [24, pp. 153–154],

$$\begin{aligned} [S(t)T(t)u_0](x) &= \tilde{S}(t) \left[\int_{-\infty}^{\infty} k^t(y) u_0(x-y) dy \right] \\ &\geq \int_{-\infty}^{\infty} k^t(y) \tilde{S}(t)[u_0(x-y)] dy \\ &= \int_{-\infty}^{\infty} k^t(y) [S(t)u_0](x-y) dy = [T(t)S(t)u_0](x). \end{aligned}$$

Thus,

$$S_N(t)T(t)u_0 = S(t)T(t)u_0 \geq T(t)S(t)u_0 = T(t)S_N(t)u_0$$

which finishes the proof by Proposition 1, (23) and (24).

Corollary 1 *Under the assumptions of Theorem 1, if u_0 and its first two derivatives in x exist and belong to $C_0(\mathbb{R})$, then (6) has a unique strong solution $u(t) = W(t)u_0$ for all $u_0 \geq 0$ in X given by the Trotter product formula (15).*

Proof In this case u_0 is in the domain of the operator A , and since the function $f(u) = ru(1 - u/K)$ as a function $X \rightarrow X$ is continuously differentiable, u is also a strong solution by [56, Chapter 6, Theorem 1.5].

Proposition 1 and Theorem 1 show that the abstract RD equation (6) can be solved by operator splitting in terms of the two component equations, the reproduction or growth equation $\dot{u} = f(u)$ and the dispersal equation $\dot{u} = Au$. This also establishes a mathematical connection between RD and ID equations. For any abstract RD Equation (1), there is a corresponding ID Equation (2) at any time scale $\tau > 0$ where $g_\tau(u) = S(\tau)u$ represents population growth over a time period of length τ , and the dispersal kernel k^τ from the infinitely divisible semi-group (13) spreads the population over the same time step. The results in this section show that solutions to this ID equation converge to the RD equation solution as the time step shrinks to zero. Hence ID equations with any infinitely divisible dispersal kernel correspond precisely to the analogous RD equation. Gaussian dispersal kernels relate to the classical Equation (1) with a Laplacian dispersal term. In the next section, we show in detail how the fractional RD Equation (3) is linked to the ID equation with a stable probability density as the dispersal kernel.

3 Fractional Reproduction-Dispersal equations

The classical diffusion equation is closely connected to the central limit theorem of statistics, since the sum of a large number of independent and statistically identical random movements converges to a normal density [24,47]. The fractional diffusion equation relates to another central limit theorem. The classical result assumes that the individual random jump has a finite standard deviation. If instead we assume that the random movements X have power-law probability tails $P(|X| > r) \approx r^{-\alpha}$ for some $0 < \alpha < 2$, then the standard deviation is infinite, and the sum converges to a stable density [24,62]. The generalised central limit theorem is completely generic, in the sense that any convergence must approach one of these forms [24]. The stable probability density functions cannot be written in closed form, except in a few special cases, so it is common to describe these distributions in terms of their Fourier transforms $\hat{k}^t(\lambda) = e^{t\psi(\lambda)}$ where

$$\psi(\lambda) = \begin{cases} -ia\lambda - \sigma^\alpha |\lambda|^\alpha (1 - i\beta(\text{sign}\lambda) \tan \frac{\pi\alpha}{2}) & \text{for } \alpha \neq 1 \\ -ia\lambda - \sigma |\lambda| (1 + i\beta \frac{2}{\pi}(\text{sign}\lambda) \ln \lambda) & \text{for } \alpha = 1. \end{cases}$$

The parameter $a \in \mathbb{R}$ centers the distribution, while $\sigma \geq 0$ provides a length scale. The stable index α and the parameter β that controls the skewness satisfy $0 < \alpha \leq 2$ and $-1 \leq \beta \leq 1$, [62]. This formula comes from computing the integral (12) with jump intensity $\phi\{x: |x| > r\} = Cr^{-\alpha}$ [47, Section 7.3]. This jump intensity comes from a regular variation argument, and reflects the power-law jumps in the limit theorem [47, Section 8.2].

Stable densities are universal dispersal kernels, since any dispersal kernel converges to one of the stable densities (including the Gaussian, the special case $\alpha = 2$) after a number of convolutions. The stochastic process with stable transition densities k^t is called a Lévy motion, a generalised form of Brownian motion that allows for occasional large jumps. A random path in this model is a fractal of dimension α [75]. The stable densities possess a pleasant scaling property $k^t(x) = t^{-1/\alpha} k(xt^{-1/\alpha})$. Since the parameter α codes the scaling, the order of the derivative, and the fractal dimension, there are several possibilities for model fitting [1–3, 9, 10, 29, 54].

The fractional diffusion equation models anomalous dispersion, the accumulation of random movements with power law probability tails [17,45]. Solutions to the fractional dispersion equation are stable probability densities [43,71]. Using the convention $(re^{i\theta})^\alpha = r^\alpha e^{i\alpha\theta}$ for $-\pi < \theta \leq \pi$, we have

$$|\lambda|^\alpha (1 - i\beta(\text{sign}\lambda) \tan \frac{\pi\alpha}{2}) = \frac{1}{\cos(\pi\alpha/2)} \left(\frac{1+\beta}{2} (i\lambda)^\alpha + \frac{1-\beta}{2} (-i\lambda)^\alpha \right).$$

Hence we can also write

$$\begin{aligned} \hat{k}^t(\lambda) &= \exp \left[t \left(-ai\lambda - \frac{\sigma^\alpha}{\cos(\pi\alpha/2)} \left(\frac{1+\beta}{2} (i\lambda)^\alpha + \frac{1-\beta}{2} (-i\lambda)^\alpha \right) \right) \right] \\ &= \exp [-vt(i\lambda) + pDt(i\lambda)^\alpha + qDt(-i\lambda)^\alpha] \end{aligned} \quad (25)$$

where $v = a$, $\sigma^\alpha = -D \cos(\pi\alpha/2)$, and $p - q = \beta$. Then the Fourier transform \hat{k}^t solves the ordinary differential equation

$$\frac{d}{dt} \hat{k}^t(\lambda) = [-v(i\lambda) + pD(i\lambda)^\alpha + qD(-i\lambda)^\alpha] \hat{k}^t(\lambda). \quad (26)$$

Recall that $(i\lambda)^n \hat{f}(\lambda)$ is the Fourier transform of $d^n f(x)/dx^n$. Similarly, we define the fractional derivative $d^\alpha f(x)/dx^\alpha$ as the function whose Fourier transform is $(i\lambda)^\alpha \hat{f}(\lambda)$. We also define $d^\alpha f(x)/d(-x)^\alpha$ as the function whose Fourier transform is $(-i\lambda)^\alpha \hat{f}(\lambda)$. Now inverting (26) yields the fractional dispersion/diffusion equation

$$\frac{\partial k^t(x)}{\partial t} = -v \frac{\partial k^t(x)}{\partial x} + pD \frac{\partial^\alpha k^t(x)}{\partial x^\alpha} + qD \frac{\partial^\alpha k^t(x)}{\partial (-x)^\alpha}. \quad (27)$$

As a stochastic model for population movements, this fractional dispersion equation differs from the classical diffusion/dispersion equation by allowing for occasional large movements many times larger than average. The resulting population density curves retain the power law tails of the individual jump distributions, and super-dispersive spreading like $t^{1/\alpha}$ makes this a useful model for populations that grow and spread faster and farther than the classical equation predicts. In the limit theorem, the weights p, q are the probability of large jumps in the positive or negative direction, respectively. Hence, the special case $\partial k^t(x)/\partial t = D \partial^\alpha k^t(x)/\partial x^\alpha$ corresponds to zero drift $v = 0$ and only positive jumps $p = 1 - q = 0$. For symmetric jumps $p = q = 1/2$ we get a symmetric dispersal kernel that solves the (Riesz) fractional dispersion equation

$$\frac{\partial k^t(x)}{\partial t} = -v \frac{\partial k^t(x)}{\partial x} + \sigma^\alpha \frac{\partial^\alpha k^t(x)}{\partial |x|^\alpha} \quad (28)$$

where $d^\alpha f(x)/d|x|^\alpha$ is defined as the function whose Fourier transform is $-|k|^\alpha \hat{f}(\lambda)$. This derivative operator is a classical fractional power of the second derivative or Laplacian [4].




Fig. 1 Standard stable dispersal kernels with $\alpha = 1.6$ illustrating the bell shape and skewness.

Solving the fractional dispersion equation with drift (27) in the constant coefficient case is equivalent to computing the stable densities. Although the stable

Fourier transform cannot be inverted in closed form, probabilists have developed fast numerical methods for computing them, based on analytical expressions obtained from the Fourier inversion formula [54]. Figure 1 illustrates the numerical calculation of the stable density using Nolan's method. Stable densities are similar to the Gaussian end-member, but can incorporate both skewness and heavy tails.

For fractional dispersion equations with variable coefficients, efficient finite difference schemes [22, 42, 73, 74] and particle tracking codes [78] have recently become available. Finite difference schemes are based on the Grünwald formula

$$\frac{d^\alpha f(x)}{dx^\alpha} = \lim_{h \rightarrow 0} \frac{1}{h^\alpha} \sum_{k=0}^{\infty} \frac{\Gamma(k-\alpha)}{k! \Gamma(-\alpha)} f(x-kh). \quad (29)$$

This limit is a Riemann-Liouville fractional derivative [57, 61]

$$\frac{d^\alpha f(x)}{dx^\alpha} = \frac{1}{\Gamma(n-\alpha)} \frac{d^n}{dx^n} \int_0^x \frac{f(y)}{(x-y)^{\alpha+1-n}} dy \quad (30)$$

where $n-1 < \alpha \leq n$. If α is an integer, then the above definitions give the standard integer derivatives. For $0 < \alpha < 1$, integration by parts in (30) yields

$$\frac{d^\alpha f(x)}{dx^\alpha} = \frac{\alpha}{\Gamma(1-\alpha)} \int_0^\infty \frac{f(x) - f(x-y)}{y} y^{-\alpha} dy \quad (31)$$

a weighted average of difference quotients, with power law weights deriving from the underlying jump intensity or Lévy measure.

Now we consider the fractional Fisher's equation

$$\frac{\partial u}{\partial t}(x,t) = ru(x,t) \left(1 - \frac{u(x,t)}{K}\right) + \sigma^\alpha \frac{\partial^\alpha u}{\partial |x|^\alpha}(x,t); \quad u(x,0) = u_0(x). \quad (32)$$

To apply the sequential operator splitting procedure from the previous section, we note that the fractional diffusion equation

$$\frac{\partial u}{\partial t}(x,t) = \sigma^\alpha \frac{\partial^\alpha u}{\partial |x|^\alpha}(x,t); \quad u(x,0) = u_0(x) \quad (33)$$

has the unique solution

$$u(x,t) = [T(t)u_0](x) = \int_{-\infty}^{\infty} k^t(x-y)u_0(y) dy \quad (34)$$

where $k^t(x)$ is a stable density with parameters $1 < \alpha \leq 2$, $\beta = 0$, $a = 0$, and $\sigma^\alpha > 0$. Next consider the growth equation

$$\frac{\partial u}{\partial t}(x,t) = ru(x,t) \left(1 - \frac{u(x,t)}{K}\right); \quad u(x,0) = u_0(x)$$

with solution (17). Define an iteration based on the sequential splitting by

$$\begin{aligned} u_{n+1}(x) &= [T(\tau)S(\tau)u_n](x) \\ &= \int_{-\infty}^{\infty} k^\tau(x-y)K \left(1 - \frac{K - u_n(y)}{K + u_n(y)(\exp(r\tau) - 1)}\right) dy. \end{aligned} \quad (35)$$

This is a discrete time ID equation of the form (5), an approximate solution $u(x, n\tau)$ of the fractional RD Equation (32). Similarly, if we model first dispersion and then growth, we obtain the alternative sequential splitting

$$\begin{aligned} u_{n+1}(x) &:= [S(\tau)T(\tau)u_n](x) \\ &= K \left(1 - \frac{K - \left(\int_{-\infty}^{\infty} k^\tau(x-y)u_n(y) dy \right)}{K + \left(\int_{-\infty}^{\infty} k^\tau(x-y)u_n(y) dy \right) (\exp(r\tau) - 1)} \right). \end{aligned} \quad (36)$$

Theorem 1 and Corollary 1 imply that the fractional Fisher's Equation (32) in continuous time can be solved numerically by computing solutions to one of its discrete time counterparts (35) or (36) with $\tau = t/n$. The approximate solutions $u_n(x, t)$ converge to the unique solution $u(x, t)$ of the fractional Fisher's equation at any time $t > 0$ for any smooth initial population density $u_0(x)$ as $n \rightarrow \infty$. This result links the continuous time partial differential equation model with the corresponding discrete time integro-difference equation model. Furthermore, the approximation (35) gives a lower bound to the exact solution while the approximation (36) gives an upper bound. Hence these two approximation can be compared to yield exact error bounds. See Section 4 for an illustration.

Theorem 1 and Corollary 1 extend immediately to any abstract RD equation of the form (6) as long as $u \mapsto [\tilde{S}(t)](u)$ is concave down on some interval $u \in [0, M]$ for each $t > 0$, and solutions to the differential equation $\dot{u} = f(u)$ remain in this interval for all time whenever $u(0) \in [0, M]$. They also extend to the case where $f(u, x)$ depends on the point x in space, and/or the multidimensional case $x \in \mathbb{R}^d$, since all the proofs are point-wise. The results of this paper also extend to model patchy populations in d -dimensional space, where the growth rate and carrying capacity vary with spatial location [7].

A wide variety of alternative models can be obtained by considering different infinitely divisible densities k as the dispersal kernel. The infinitely divisible densities are convenient because one can define the integro-difference Equation (5) at any time scale τ based on the convolution power k^τ . The results of this paper also yield the corresponding RD model with a diffusion operator A exhibiting as the generator of the corresponding infinitely divisible semigroup. Lockwood et al. [38] employ a Laplace (double exponential), or more generally, a double Gamma family of dispersion kernels, which are infinitely divisible with Lévy representation $[a, 0, \phi]$ with jump intensity $\phi(y) = |y|^{-1} e^{-c|y|}$. In this case the last term in the integral (14) converges, and hence we can choose a so that

$$Au(x) = \int_{-\infty}^{\infty} \frac{u(x-y) - u(x)}{|y|} e^{-c|y|} dy \quad (37)$$

an *exponentially weighted derivative*, which is similar to the fractional derivative formula (31) but with different weights [37]. For an exponential or Gamma dispersal kernel, the generator formula is the same as (37) except that the integral is taken over $y > 0$. Clarke et al. [18, 19] employ the Student- t dispersal kernel, which is infinitely divisible with Lévy representation as specified in [28, Remark 2.3] in terms of Bessel functions. Substituting into (14) yields the corresponding

generator, connecting the integro-difference equation (5) with a Student- t dispersal kernel to the analogous reaction-diffusion Equation (4).

The fractional diffusion equation is a mathematical abstraction of the real situation in which a power law dispersal kernel pertains at some scale. Realistically, the power law probability density should not extend to infinity, since this would imply an unbounded velocity distribution. Of course, the same observation holds for the Gaussian, or any other dispersal kernel that extends to infinity. In the Gaussian case, the central limit theorem implies that the additive effect of a number of independent movements will approach a Gaussian form. For power law kernels, the extended central limit theorem implies an approach to a stable kernel. Truncated Lévy flights [39, 70] and the corresponding fractional diffusion equation [72] provide a more realistic model that imposes a cutoff on the power-law tail. While the asymptotics are Gaussian at a sufficiently long time scale, the stable fit excels at the field scale [40, 41, 76], and thus provides the simplest useful approximation.

4 Numerical Experiments

Consider a fractional Fisher's equation and a symmetric (Riesz) fractional diffusion term of order $1 \leq \alpha \leq 2$:

$$\frac{\partial u}{\partial t}(x, t) = \sigma^\alpha \frac{\partial^\alpha u}{\partial |x|^\alpha}(x, t) + ru(x, t) \left(1 - \frac{u(x, t)}{K}\right), \quad u(x, 0) = u_0(x). \quad (38)$$

Solutions to this equation can be computed using the results of Theorem 1, using the integro-difference model (5) as an approximation where the dispersal kernel $k^\tau(y)$ is a symmetric α -stable probability density function with index α , skewness $\beta = 0$, centre $a = 0$, and scale $\tau\sigma$. Note that in case $\alpha = 1$, k^τ is the density of a Cauchy distribution as used for example in [36, 69].




Fig. 2 Solution of the fractional Fisher's equation (38) with $\alpha = 1.8$ versus $\alpha = 2$ at $t = 32$ with $K = 1$, $r = 0.25$ and $\sigma^\alpha = 0.1$ showing heavier tails and faster spreading in the fractional case $\alpha < 2$.

Numerical simulations were performed with $K = 1$, $c = 0.25$ and $\sigma^\alpha = 0.1$ and a smooth step-like initial function u_0 which takes the constant value $u = 0.8$

around the origin and rapidly decays to 0 away from the origin. $S(\tau)$ was computed analytically at each time step using the explicit analytical solution (17) and $T(\tau)u_n$ was computed by numerically convolving u_n against the symmetric α -stable dispersal kernel k^τ , which was computed numerically using the method of Nolan [54]. Note that the dispersal kernel can be computed for any time scale τ , and that it only has to be computed once for each time scale. Figure 2 shows that the use of the conventional diffusion term $\alpha = 2$ in (38) produces a rapidly decaying solution away from the origin. However, if one replaces the diffusion term by a fractional one, even with an order α which is close to 2, then the solution picks up heavy tails and is spreading faster, similar to the results reported in del-Castillo-Negrete et al. [21] for the one-sided fractional RD Equation (3).




Fig. 3 Sequential splitting approximations of the solution of the fractional Fisher's Equation (38) with $\alpha = 1.8$, $K = 1$, $r = 0.25$ and $\sigma^\alpha = 0.1$. Dashed lines are computed from (36) with time step $\tau = 32, 8, 2$ and dotted lines are computed from (35) at the same time steps to show monotone convergence to the exact (solid line) solution.

Figure 3 visualises the conclusions of Theorem 1; that is, the sequential splitting approximations (35) and (36) converge in a pointwise monotone increasing and decreasing fashion, respectively, to the solution of (38).

Figure 4 shows evidence of an accelerating front. The selected population density level $u = c$ has spread a distance $x_c(t)$ by time $t > 0$, and since the graph of $x_c(t)$ versus t for various c closely resembles a straight line on the semi-log plot, we conclude that the front expands exponentially with time, in agreement with results reported by del-Castillo-Negrete et al. [21] for the one-sided model (3).

5 Conclusions

We showed how any integro-difference equation with an infinitely divisible dispersal kernel can be approximated by an integro-differential equation. Amongst




Fig. 4 Exponential spreading front for the fractional Fisher's equation (38) with $\alpha = 1.8$, $K = 1$, $r = 0.25$ and $\sigma^\alpha = 0.1$. The farthest distance to which a population density of c has spread by time t is plotted on a semi-log scale to illustrate the (nearly) exponential front growth over time.

the many infinitely divisible kernels there are only a few that are densities of stable probability distributions. They are the scaling limit of sums of independent identically distributed random variables and keep their shape at any time scale; the most popular being the Gaussian distribution. The classical central limit theorem implies that the Gaussian is the scaling limit of any distribution that does not have a power law tail. In case the dispersal kernels are power law of index $\alpha < 2$, the limit will be a stable distribution of index α . These stable distributions in general have no closed form description in real space (only Cauchy, Levy, and Gaussian), but are easily described in Fourier space. In the same way a Gaussian kernel gives rise to a Laplace operator in the differential equation, stable kernels give rise to fractional derivative operators and hence similarly parsimonious models. We investigated these fractional models, showed that solutions can be well approximated numerically and that they give rise to an accelerating invasion front.

6 Acknowledgements

We would like to thank Shona Lamoureaux of AgResearch New Zealand, Jon Pitchford at the University of Leeds, and Alex James at the University of Canterbury, New Zealand for stimulating discussions.

References

1. Aban, I.B., Meerschaert, M. M., 2001. Shifted Hill's estimator for heavy tails, *Communications in Statistics: Simulation and Computation* 30(4), 949–962.
2. Aban, I.B., Meerschaert, M.M., 2004. Generalized least squares estimators for the thickness of heavy tails, *Journal of Statistical Planning and Inference* 119(2), 341–352.

3. Aban, I.B., Meerschaert, M.M., Panorska, A.K., 2006. Parameter estimation for the truncated Pareto distribution, *Journal of the American Statistical Association: Theory and Methods*, 101(473), 270–277.
4. Arendt, W., Batty, C., Hieber, M., Neubrander, F. 2001. *Vector-valued Laplace transforms and Cauchy problems*, Monographs in Mathematics. Birkhaeuser-Verlag, Berlin.
5. Arendt, W., Grabosch, A., Greiner, G., Groh, U., Lotz, H.P., Moustakas, U., Nagel, R., Neubrander, F., Schlotterbeck, U. 1986. *One-parameter semigroups of positive operators*, Lecture Notes in Mathematics, vol. 1184. Springer-Verlag, Berlin.
6. Baeumer, B., Meerschaert, M.M., Benson, D.A., Wheatcraft, S.W., 2001. Subordinated advection-dispersion equation for contaminant transport, *Water Resources Research* 37, 1543–1550.
7. Baeumer, B., Meerschaert, M.M., Kovacs, M., to appear. Numerical solutions for fractional reaction-diffusion equations, *Comput. Math. Appl.*
8. Barkai, E., Metzler, R., Klafter, J., 2000. From continuous time random walks to the fractional Fokker-Planck equation, *Phys. Rev. E* 61, 132–138.
9. Benson, D., Wheatcraft, S., Meerschaert, M., 2000. Application of a fractional advection-dispersion equation, *Water Resources Research* 36(6), 1403–1412.
10. Benson, D. A., Schumer, R., Meerschaert, M. M., Wheatcraft, S.W., 2001. Fractional dispersion, Lévy motions, and the MADE tracer tests, *Transport in Porous Media* 42, 211–240.
11. Benson, D. A., Wheatcraft, S. W., Meerschaert, M. M., 2000. The fractional-order governing equation of Lévy motion, *Water Resources Research* 36, 1413–1424.
12. Blumen, A., Zumofen, G., Klafter, J., 1989. Transport aspects in anomalous diffusion: Lévy walks, *Phys. Rev. A* 40, 3964–3973.
13. Bouchaud, J., Georges, A., 1990. Anomalous diffusion in disordered media - statistical mechanisms, models and physical applications, *Phys. Rep.* 195, 127–293.
14. Brézis, H., Pazy, A., 1972. Convergence and approximation of semigroups of nonlinear operators in Banach spaces, *J. Functional Analysis* 9, 63–74.
15. Bullock, J.M., Clarke, R.T., 2000. Long distance seed dispersal by wind: measuring and modelling the tail of the curve, *Oecologia* 124(4), 506–521.
16. Chambers, J.M., Mallows, C.L., Stuck, B.W., 1976. A method for simulating stable random variables, *J. Amer. Statist. Assoc.* 71(354), 340–344.
17. Chaves, A., 1998. A fractional diffusion equation to describe Lévy flights, *Phys. Lett. A* 239, 13–16.
18. Clark, J.S., Silman, M., Kern, R., Macklin, E., HilleRisLambers, J., 1999. Seed dispersal near and far: Patterns across temperate and tropical forests, *Ecology* 80(5), 1475–1494.
19. Clark, J. S., Lewis, M., Horvath, L., 2001. Invasion by Extremes: Population Spread with Variation in Dispersal and Reproduction, *The American Naturalist* 157(5), 537–554.
20. Cliff, M., Goldstein, J. A., Wacker, M., 2004. Positivity, Trotter products, and blow-up, *Positivity* 8(2), 187–208.
21. del Castillo-Negrete, D., Carreras, B.A., Lynch, V.E., 2003. Front dynamics in reaction-diffusion systems with levy flights: A fractional diffusion approach, *Physical Review Letters* 91(1), 018302.
22. Deng, Z., Singh, V. P., Bengtsson L., 2004. Numerical solution of fractional advection-dispersion equation, *Journal of Hydraulic Engineering* 130, 422–431.
23. Engel, K.J., Nagel, R., 2000. *One-parameter semigroups for linear evolution equations*, Graduate Texts in Mathematics, vol. 194. Springer-Verlag, New York.
24. Feller, W., 1966. *An Introduction to Probability Theory and Applications*. Volumes I and II, John Wiley and Sons.
25. Fisher, R.A., 1937. The wave of advances of advantageous genes, *Annals of Eugenics* 7, 353–369.
26. Gnedenko, B., Kolmogorov, A., 1968. *Limit Distributions for Sums of Independent Random Variables*. Translated from the Russian, annotated, and revised by K. L. Chung. With appendices by J. L. Doob and P. L. Hsu. Revised edition. Addison-Wesley, Reading, Mass.
27. Gorenflo, R., Mainardi, F., Scalas, E., Raberto, M., 2001. Problems with using existing transport models to describe microbial transport in porous media, In: *Trends Math.*, pp. 171–180. American Society for Microbiology, Mathematical finance (Konstanz, 2000).
28. Heyde, C.C., Leonenko, N.N., 2005. Student Processes, *Advances in Applied Probability* 37, 342–365.
29. Hill, B., 1975. A simple general approach to inference about the tail of a distribution, *Ann. Statist.* 3(5), 1163–1173.

30. Hille, E., Phillips, R.S., 1974. Functional analysis and semi-groups. American Mathematical Society, Providence, R. I. Third printing of the revised edition of 1957, American Mathematical Society Colloquium Publications, Vol. XXXI.
31. Jacob, N., 1996. Pseudo-differential operators and Markov processes, Mathematical Research, vol. 94. Akademie Verlag, Berlin.
32. Katul, G. G., Porporato, A., Nathan, R., Siqueira, M., Soons, M. B., Poggi, D., Horn, H. S., Levin, S.A. 2005. Mechanistic analytical models for long-distance seed dispersal by wind, *The American Naturalist* 166, 368–381.
33. Klafter, J., Blumen, A., Shlesinger, M.F., 1987. Stochastic pathways to anomalous diffusion, *Phys. Rev. A* 35, 3081–3085.
34. Klein, E. K., Lavigne, C., Picault, H., Renard, M., Gouyon, P. H., 2006. Pollen dispersal of oilseed rape: estimation of the dispersal function and effects of field dimension, *Journal of Applied Ecology* 43(10), 141–151.
35. Kolmogorov, A., Petrovsky, N., Piscounov, S., 1937. Étude de l'équations de la diffusion avec croissance de la quantité de matière et son application a un problème biologique, *Moscow University Bulletin of Mathematics* 1, 1–25.
36. Kot, M., Lewis, M. A., Van den Driessche, P., 1996. Dispersal data and the spread of invading organisms, *Ecology* 77, 2027–2042.
37. Kozubowski, T., Meerschaert, M., Podgórski, K., 2006. Fractional Laplace motion, *Advances in Applied Probability* 38(2), 451–464.
38. Lockwood, D. R., Hastings, A., 2002. The effects of dispersal patterns on marine reserves: Does the tail wag the dog? *Theoretical Population Biology* 61, 297–309.
39. Mantegna, R. N., Stanley, H. E., 1994. Stochastic process with ultraslow convergence to a Gaussian: The truncated Lévy flight, *Phys. Rev. Lett.* 73, 2946.
40. Mantegna, R. N., Stanley, H. E., 1997. Econophysics: Scaling and its Breakdown in Finance, *J. Stat. Phys.* 89, 469–479.
41. Mantegna, R. N., Stanley, H. E., 1998. Modeling of financial data: Comparison of the truncated Lévy flight and the ARCH(1) and GARCH(1,1) processes, *Physica A* 254(1), 77–84.
42. Lynch, V. E., Carreras, B. A., del-Castillo-Negrete, D., Ferreira-Mejias, K. M., Hicks, H. R., 2003. Numerical methods for the solution of partial differential equations of fractional order, *J. Comput. Phys.* 192, 406–421.
43. Meerschaert, M.M., Benson, D.A., Baeumer, B., 1999. Multidimensional advection and fractional dispersion, *Phys. Rev. E* 59, 5026–5028.
44. Meerschaert, M.M., Benson, D.A., Baeumer, B., 2001. Operator Lévy motion and multi-scaling anomalous diffusion. *Phys. Rev. E* 63, 1112–1117.
45. Meerschaert, M.M., Benson, D.A., Scheffler, H., Baeumer, B., 2002. Stochastic solution of space-time fractional diffusion equations, *Phys. Rev. E* 65, 1103–1106.
46. Meerschaert, M.M., Benson, D.A., Scheffler, H.P., Becker-Kern, P., 2002. Governing equations and solutions of anomalous random walk limits, *Phys. Rev. E* 66, 102R–105R.
47. Meerschaert, M.M., Scheffler, H.P., 2001. *Limit Distributions for Sums of Independent Random Vectors: Heavy Tails in Theory and Practice*, Wiley Interscience, New York.
48. Metzler, R., Klafter, J., 2000. The random walk's guide to anomalous diffusion: A fractional dynamics approach. *Phys. Rep.* 339, 1–77.
49. Metzler, R., Klafter, J., 2004. The restaurant at the end of the random walk: recent developments in the description of anomalous transport by fractional dynamics. *J. Physics A* 37, R161–R208.
50. Miller, K., Ross, B., 1993. *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley and Sons, New York.
51. Miyadera, I., Ôharu, S., 1970. Approximation of semi-groups of nonlinear operators, *Tôhoku Math. J.* 22, 24–47.
52. Murray, J.D., 2002. *Mathematical biology. I,II, Interdisciplinary Applied Mathematics*, vol. 17,18, third edn., Springer-Verlag, New York.
53. Neubert, M., Caswell, H., 2000. Demography and dispersal: Calculation and sensitivity analysis of invasion speed for structured populations, *Ecology* 81(6), 1613–1628.
54. Nolan, J.P., 1997. Numerical calculation of stable densities and distribution functions. Heavy tails and highly volatile phenomena, *Comm. Statist. Stochastic Models* 13(4), 759–774.
55. Paradis, E., Baillie, S.R., Sutherland, W.J., 2002. Modeling large-scale dispersal distances. *Ecological Modelling* 151(2-3), 279–292.

-
56. Pazy, A., 1983. *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Applied Mathematical Sciences, vol. 44. Springer-Verlag, New York.
 57. Podlubny, I., 1999. *Fractional Differential Equations*. Academic Press, New York.
 58. Raberto, M., Scalas, E., Mainardi, F., 2002. Waiting-times and returns in high-frequency financial data: an empirical study, *Physica A* 314, 749–755.
 59. Sabatelli, L., Keating, S., Dudley, J., Richmond, P., 2002. Waiting time distributions in financial markets, *Eur. Phys. J. B* 27, 273–275.
 60. Saichev, A.I., Zaslavsky, G.M., 1997. Fractional kinetic equations: solutions and applications, *Chaos* 7(4), 753–764.
 61. Samko, S., Kilbas, A., Marichev, O., 1993. *Fractional Integrals and derivatives: Theory and Applications*, Gordon and Breach, London.
 62. Samorodnitsky, G., Taqqu, M.S., 1994. *Stable Non-Gaussian Random Processes*, Chapman & Hall/CRC.
 63. Sato, K., 1999. *Lévy Processes and Infinitely Divisible Distributions*, Cambridge studies in advanced mathematics, Cambridge University Press.
 64. Scalas, E., Gorenflo, R., Mainardi, F., 2000. Fractional calculus and continuous-time finance, *Phys. A* 284, 376–384.
 65. Scalas, E., 2006. The application of continuous-time random walks in finance and economics, *Physica A* 362, 225–239.
 66. Scalas, E., Gorenflo, R., Luckcock, H., Mainardi, F., Mantelli, M., Raberto, M., 2005. On the Intertrade Waiting-time Distribution, *Finance Letters* 3, 38–43.
 67. Schumer, R., Benson, D. A., Meerschaert, M. M., Baeumer, B., 2003. Multiscaling fractional advection-dispersion equations and their solutions, *Water Resources Research* 39, 1022–1032.
 68. Schumer, R., Benson, D. A., Meerschaert, M. M., Wheatcraft, S. W., 2001. Eulerian derivation of the fractional advection-dispersion equation, *Journal of Contaminant Hydrology* 48, 69–88.
 69. Shaw M.W., 1995. Simulation of population expansion and spatial pattern when individual dispersal distributions do not decline exponentially with distance. *Proceedings of the Royal Society of London Series B* 259, 243–248.
 70. Shlesinger, M. F., 1995. Comment on Stochastic Process with Ultraslow Convergence to a Gaussian: The Truncated Lévy Flight, *Phys. Rev. Lett.* 74, 4959–4959.
 71. Sokolov, I. M., Klafter, J., 2005. From diffusion to anomalous diffusion: a century after Einstein’s Brownian motion, *Chaos* 15(2), 026103-1–026103-7.
 72. Sokolov, I. M., Chechkin, A. V., Klafter, J., 2004. Fractional diffusion equation for a power-law-truncated Lévy process, *Physica A* 336(3–4), 245–251.
 73. Tadjeran, C., Meerschaert, M. M., Scheffler, H. P., 2006. A second order accurate numerical approximation for the fractional diffusion equation, *J. Comput. Phys.* 213, 205–213.
 74. Tadjeran, C., Meerschaert, M. M., 2007. A second order accurate numerical method for the two-dimensional fractional diffusion equation, *J. Comput. Phys.* 220 pp. 813823.
 75. Taylor, S.J., 1986. The measure theory of random fractals, *Math. Proc. Cambridge Philos. Soc.* 100, 383–406.
 76. Viswanathan, G. M., Afanasyev, V., Buldyrev, S. V., Murphy, E. J., Prince, P. A., Stanley, H. E., 1996. Lévy flight search patterns of wandering albatrosses, *Nature* 381, 413–415.
 77. Zaslavsky, G., 1994. Fractional kinetic equation for hamiltonian chaos. Chaotic advection, tracer dynamics and turbulent dispersion, *Phys. D* 76, 110–122.
 78. Zhang, Y., Benson, D. A., Meerschaert, M. M., Scheffler, H. P., 2006. On using random walks to solve the space-fractional advection-dispersion equations, *J. Statist. Phys.* 123, 89–110.